

NIH RELAIS Document Delivery

NIH-10092956

NIH -- W1 PE175

PAMELA GEHRON ROBEY
CSDB/NIDR/NIH Bldng 30 Rm 228
30 CONVENT DRIVE MSC 4320
BETHESDA, MD 20892

ATTN:	SUBMITTED:	2001-12-17 15:18:34
PHONE: 301-496-4563	PRINTED:	2001-12-18 12:19:16
FAX: 301-402-0824	REQUEST NO.:	NIH-10092956
E-MAIL:	SENT VIA:	LOAN DOC 5315875

NIH	Fiche to Paper	Journal
TITLE:	PEDIATRIC RESEARCH	
PUBLISHER/PLACE:	Lippincott Williams _Wilkins Hagerstown, Md :	
VOLUME/ISSUE/PAGES:	1986 Sep;20(9):859-63 859-63	
DATE:	1986	
AUTHOR OF ARTICLE:	Foster CM; Feuillan P; Padmanabhan V; Pescovitz OH; Beitins	
TITLE OF ARTICLE:	Ovarian function in girls with McCune-Albright syn	
ISSN:	0031-3998	
OTHER NOS/LETTERS:	Library reports holding volume or year 0100714 3092175	
SOURCE:	PubMed	
CALL NUMBER:	W1 PE175	
REQUESTER INFO:	AB424	
DELIVERY:	E-mail: probey@DIR.NIDCR.NIH.GOV	
REPLY:	Mail:	

NOTICE: THIS MATERIAL MAY BE PROTECTED BY COPYRIGHT LAW (TITLE 17, U.S. CODE)

-----National-Institutes-of-Health,-Bethesda,-MD-----

Ovarian Function in Girls with McCune-Albright Syndrome¹

CAROL M. FOSTER, PENELOPE FEUILLAN, VASANTHA PADMANABHAN, ORA H. PESCOVITZ,
INESE Z. BEITINS, FLORENCE COMITE, THOMAS H. SHAWKER, D. LYNN LORIAUX, AND
GORDON B. CUTLER, JR.

*Developmental Endocrinology Branch, National Institute of Child Health and Human Development,
Bethesda, Maryland 20892 [C.M.F., P.F., O.H.P., F.C., D.L.L., G.B.C.]; Department of Pediatrics, University of
Michigan Medical Center, Ann Arbor, Michigan 48109 [C.M.F., V.P., I.Z.B.]; and Diagnostic Radiology,
Clinical Center, National Institutes of Health, Bethesda, Maryland 20892 [T.H.S.]*

ABSTRACT. We measured plasma estradiol levels and ovarian volumes in eight girls with precocious puberty due to McCune-Albright syndrome. Six girls had gonadotropin-independent ovarian estrogen secretion and two girls had pubertal gonadotropin levels. Mean ovarian volume in all patients was significantly greater than in normal prepubertal girls. Mean ovarian volumes of the girls with McCune-Albright syndrome overlapped the range found in girls with idiopathic central precocious puberty or central precocious puberty associated with central nervous system lesions. However, the degree of asymmetry between the right and left ovaries was significantly greater in girls with McCune-Albright syndrome. Asymmetry was due, for the most part, to the presence of large solitary cysts in the larger of the two ovaries. In the six girls with McCune-Albright syndrome and gonadotropin-independent precocious puberty, both mean ovarian volume and the degree of asymmetry between the right and left ovaries were significantly correlated with plasma estradiol. Serum follicle-stimulating hormone bioactivity was increased in two patients but did not vary with ovarian cyst size. Thyroid-stimulating hormone levels were normal but serum prolactin was slightly elevated in one of the six girls with gonadotropin-independent precocious puberty. Fluctuation in the size of unilateral ovarian cysts appears to result in changes in the plasma estradiol level, leading to advancement and spontaneous regression of secondary sexual characteristics and menses in girls with McCune-Albright syndrome. The cause of the cyst formation is unknown but may be related to periodic elevation of as yet undefined serum factors such as follicle-stimulating hormone bioactive substances. (*Pediatr Res* 20: 859-863, 1986)

Abbreviations

FSH, follicle-stimulating hormone
TSH, thyroid-stimulating hormone
LH, luteinizing hormone
LHRH, luteinizing hormone-releasing hormone
CNS, central nervous system
hCG, human chorionic gonadotropin

Girls with McCune-Albright syndrome have fibrous dysplasia of bone associated with café-au-lait skin pigmentation and/or precocious puberty (1, 2). Advancement and regression of secondary sexual characteristics and episodes of menses in these girls often follow a cyclical course (3, 4) and in most girls, are independent of pubertal secretion of gonadotropins (3-6). In fact, gonadotropin levels are frequently suppressed below prepubertal levels (5).

Regression of secondary sexual characteristics in girls with McCune-Albright syndrome has been observed when ovarian cysts or wedge sections of ovaries have been removed (7, 8). Comite *et al.* (4) observed that in one girl with McCune-Albright syndrome, fluctuation of the size of a left ovarian cyst was associated with increases and decreases of plasma estradiol levels. In order to explore this association further, we compared ovarian volume and estradiol levels in eight girls who had precocious puberty associated with McCune-Albright syndrome. We have attempted to determine the etiology of cyst formation by examining serum FSH bioactivity and TSH and prolactin concentrations.

METHODS

Subjects. The initial clinical and laboratory findings of the eight girls with McCune-Albright syndrome are indicated in Table 1. All girls had fibrous dysplasia of bone and precocious puberty. Patients 1, 2, 4, and 6 had café-au-lait skin pigmentation. Patients 1 and 6 had extensive lower limb deformities with subsequent height loss. Patients 1 through 6 had prepubertal or suppressed gonadotropin levels whereas patients 7 and 8 had pubertal nocturnal LH pulsations and a pubertal gonadotropin response to injection of LHRH. Neoplasms were not detected by computed tomography of the head or by ultrasonography of the pelvis and adrenal glands. Plasma 17-hydroxyprogesterone and 11-deoxycortisol, measured to exclude congenital adrenal hyperplasia, were normal. Serum T₄ levels were also normal.

Girls with central precocious puberty, either idiopathic ($n = 33$) or associated with a CNS lesion ($n = 8$), ranged from 1 to 18 yr of age. All had stage II or greater pubertal development [according to the method of Tanner (9)], advanced bone age for chronological age, and accelerated height velocity. All had a pubertal response of gonadotropins in response to LHRH and/or nocturnal pulsations of LH. Girls were included for study only after exclusion of ovarian or adrenal neoplasms by ultrasonography, and all had normal serum 17-hydroxyprogesterone and 11-deoxycortisol.

Protocol. Patients were admitted to the Clinical Center of the National Institutes of Health. The protocol was approved by the Clinical Research Committee of the National Institute of Child

Received February 24, 1986; accepted April 25, 1986.

Address correspondence to Carol M. Foster, M.D., Department of Pediatrics, University of Michigan, C. S. Mott Hospital F2125, Ann Arbor, MI 48109.

¹ Presented in part before the Society for Pediatric Research, San Francisco, CA, May 1984.

Table 1. Clinical and hormonal characteristics*

Patient	Age (yr)	Ht (cm)	%	Pubertal stage†				Peak response to LHRH		
				Bone age (yr)	Breast	Pubic hair	Menses	LH (mIU/ml)	FSH (mIU/ml)	Estradiol (pg/ml)
1	4.5	107	60	7.8	III	IV	+	2.8	0.3	21 ± 2
2	3.2	108	97	7.8	IV	III	+	2.2	0.7	132 ± 16
3	1.2	82	95	2.0	III	I	+	1.6	0.2	133 ± 22
4	5.1	128	97	8.8	IV	III	+	2.5	10.7	<20
5	4.5	117	97	6.8	IV	II	+	7.0	9.6	<20
6	5.2	102	25	6.0	II	II	+	2.0	5.7	<20
7	10.5	141	50	12.0	IV	II	+	99.8	20.0	45 ± 6
8	8.9	152	97	13.5	IV	III	+	123	22.8	50 ± 4

* Data for patients 1, 3 to 6, and 8 has been shown previously (3).

† According to the method of Tanner (9).

Health and Human Development. Informed consent was obtained from a parent and assent was obtained from children more than 7 yr of age.

LHRH stimulation was performed by administration of 100 µg of LHRH intravenous at time 0 and gonadotropin levels were determined at -30, -15, 0, 15, 30, 45, 60, 90, and 120 min. Estradiol was determined as the mean of four values obtained at 1000, 1400, 2200, and 0200 h. Children with McCune-Albright syndrome were either followed without treatment or treated with the long-acting agonist of LHRH, D-Trp⁶-Pro⁹-NEt-LHRH (LHRH_a) at a dose of 4 µg/kg/day subcutaneous. This analog has been shown to suppress gonadotropins thereby suppressing estradiol levels in girls with central precocious puberty (10, 11), but has not influenced estradiol levels in girls with gonadotropin-independent ovarian estrogen secretion (4, 5). Serial ultrasounds and estradiol measurements (mean of four values as above) were obtained in patients 1, 2, 3, and 5 at approximately 3-month intervals.

Hormone assays. Serum LH, FSH, and estradiol were measured by modifications of previously described methods (Second International Reference Preparation of Human Menopausal gonadotropin) (12-14). Sensitivity of the estradiol assay was 20 pg/ml, and intra- and interassay coefficients of variation were 8 and 16%, respectively. TSH and prolactin were measured by radioimmunoassay as previously described (15, 16).

FSH bioassay. FSH bioactivity was quantified by means of FSH-dependent aromatase activity (conversion of androgen substrate to estradiol). Following decapitation, testes from 7- to 10-day-old Sprague Dawley rats were decapsulated and incubated in a combination of Dulbecco's and Ham's F10 (1:1) medium containing 0.03% collagenase and 0.003% soybean trypsin inhibitor for 5-10 min at 34° C. Medium was decanted to remove interstitial cells. The tubules were washed and resuspended in medium containing 0.03% collagenase, 0.003% soy trypsin inhibitor, and 0.03% DNase for 30 min at 34° C. Dispersion of Sertoli cells was hastened by mechanical dispersion with a pasteur pipette. The resulting cell suspension was washed three times, and cells resuspended in medium [formulation based on Rich *et al.* (17)] containing 19-OH androstenedione (2.5 µM) and methyl isobutyl xanthine (0.1 mM). One ml of cell suspension was transferred to each well of a multiwell culture dish (Falcon, diameter 16 mm). The cells were washed and reincubated in incubation medium containing increasing concentrations NIAMDD hFSH 1-2 (AFP 2844B, biopotency 3925 IU/mg; immunopotency 5226 IU/mg) or with unknown serum samples (3 and 5 µl). Following a 24-h incubation, medium was aspirated and assayed for estradiol by radioimmunoassay (18). The sensitivity of the bioassay system was 0.01 ng/ml and index of precision 0.049. The assay is specific for FSH bioactivity (19) (Padmanabhan V, unpublished data).

Ultrasonography. Pelvic ultrasonography, using a static B-scanner, duplicated that described by Sample *et al.* (20). Ovarian

volumes were calculated by the 3-dimensional diameters from sagittal and transverse scanning and then using the formula: volume = 0.5 (width × thickness × length).

Statistical analysis. Data are represented as the mean ± SEM. Statistical comparisons were made by Student's *t* test. Correlations of estradiol and ovarian volume were determined by least squares linear regression analysis.

RESULTS

Ovarian volumes in McCune-Albright syndrome and in central precocious puberty. Mean volumes in girls with McCune-Albright syndrome overlapped the range seen in girls with idiopathic or CNS lesion associated central precocious puberty (Fig. 1A). However, when individual ovarian volumes were compared, girls with McCune-Albright syndrome had greater asymmetry of ovarian volume than did the girls with central precocious puberty (Fig. 1B). The difference in right to left ovarian volume was significantly greater in girls with McCune-Albright syndrome ($p < 0.05$, difference of 5.5 ± 1.5 ml) as compared to those with central precocious puberty (1.5 ± 0.2 ml). In patients 2, 3, and 4, a cyst of greater than 1 cm in diameter was present in the larger ovary. In patients 2 through 6, large cysts were documented at various times, unilaterally, during longitudinal study. The cysts always occurred in the same ovary in patients 3, 4, and 5. Cysts were documented in both ovaries of patient 2 although asymmetry persisted and the right ovary was always larger than the left when estradiol was elevated. In patient 6, the left ovary had been previously excised because of the presence of a large ovarian cyst. Subsequently, the right ovary developed a large cyst. In patient 8, multiple small cysts were noted in both ovaries. Cysts were not seen in ultrasounds of patients 1 and 7.

Estradiol levels and ovarian volume. The girls with McCune-Albright syndrome initially presented with estradiol levels ranging from less than 20 to 133 pg/ml although all had evidence of advanced secondary sexual characteristics and advancement of bone age. This indicated that estradiol levels must have been elevated in all these girls and may have fluctuated over time. We compared the ovarian volumes of the girls with undetectable estradiol levels at their initial visit (less than 20 pg/ml) to those with detectable estradiol levels (greater than 20 pg/ml). The girls with detectable levels had significantly larger (7.6 ± 1.2 ml) ovarian volumes than did girls with undetectable estradiol levels (2.8 ± 0.8 ml, $p < 0.02$). Patients 1, 2, 3, and 5, who had low gonadotropin levels, were followed longitudinally for 1 or more yr at 3- to 6-month intervals with pelvic ultrasound examinations and determination of estradiol levels. These girls had been treated for 3-12 months with LHRH_a but had shown no decrease in serum gonadotropin levels nor in estradiol fluctuation (5). Over time, mean ovarian volumes fluctuated and were increased in association with elevation of estradiol levels [$r = 0.79$, $p < 0.001$ (Fig. 2A)]. Similarly, the differences in right to left ovarian

MEAN OVARIAN VOLUME (ml)

OVARIAN VOLUME (ml)

Fig. 1. Compared precocious puberty, et al. (29) present anterior) area repre ml). B. di

volume = 0.72, p Serum occasions bleeding of precoc




Fig. 1. Ovarian volume of girls with McCune-Albright syndrome as compared the ovarian volume of girls with central true precocious puberty, either idiopathic or associated with a CNS lesion from Shawker *et al.* (29). Measurements were made by ultrasonography at initial presentation. Volume was determined as $0.5 \times (\text{transverse} \times \text{sagittal} \times \text{anterior})$ diameters of each ovary. A, mean ovarian volumes. The shaded area represents the mean ovarian volume of prepubertal subjects (<0.9 ml). B, difference between right and left ovarian volumes.

volume also correlated with increases in plasma estradiol level [$r = 0.72$, $p < 0.001$ (Fig. 2B)].

Serum prolactin and TSH levels. Girls with hypothyroidism occasionally develop secondary sexual characteristics and vaginal bleeding in association with large ovarian cysts (20). The cause of precocious puberty in girls with hypothyroidism is unknown




Fig. 2. Plasma estradiol levels and ovarian volume in girls with gonadotropin-independent precocious puberty. Data from patients 1, 2, 3, and 5, in whom sequential ultrasounds were done, are represented. Patients were either untreated or treated with LHRH_a. Estradiol is expressed as the mean of four samples obtained at 1000, 1400, 2200, and 0200. A, comparison of plasma estradiol with mean ovarian volume. B, comparison of plasma estradiol with difference between the volumes of the right and left ovaries.

but believed to be related to elevated TSH levels which may, in turn, increase gonadotropin or prolactin secretion. Also, the gonadotropins and TSH share a common α -subunit. This potentially might lead to hormonal overlap increasing gonadal activity which would result in precocious puberty. Pituitary hyperfunction has been reported in McCune-Albright syndrome especially for prolactin and growth hormone (22, 23). Thus, we examined TSH and prolactin in the six patients with gonadotropin independent precocious puberty (Table 2). The TSH levels were

Table 2. Prolactin and TSH levels

Patient	Prolactin (ng/ml)	TSH (μ U/ml)
1	11.4	2.4
2	6.2	3.5
3	36.2	4.0
4	NA*	<2.0
5	3.5	3.2
6	18.9	1.6
Normal range	<35	<6.0

* Not available.

Table 3. FSH bioactivity and ovarian volume

Patient	Ovarian volume (ml)		FSH bioactivity (ng/ml)
	Right	Left	
1	7.6	1.6	26.0
2	3.0	1.5	<4.0
	6.1	2.9	4.0
	39.6	9.6	2.0
3	1.0	13.0	8.0
	0.8	3.1	4.2
	1.8	3.2	<4.0
5	2.6	0.9	22.0
	16.8	4.0	20.0
	1.9	0.8	<4.0
Prepubertal normals	<0.9*	<0.9*	†

* Based on normals by Sample *et al.* (19).

† FSH bioactivity has not been detected in normal prepubertal subjects to date.

normal in all subjects tested. Prolactin was slightly increased in one subject at the time she had a large left ovary (13 ml) with a cyst. However, prolactin was normal in the remaining four subjects tested regardless of ovarian size. None of the patients exhibited hyperthyroidism, as assessed by T_4 radioimmunoassay at the time of study.

Serum FSH bioactivity. It is believed that complete ovarian function cannot occur without FSH priming, principally since girls with hCG-producing tumors have not been reported to develop precocious puberty. Our patients with McCune-Albright syndrome appeared to have functioning ovarian cysts but low to suppressed levels of FSH. It is possible that the serum of these girls might have a substance with FSH bioactivity that was not recognized in our FSH radioimmunoassay. FSH bioactivity was determined in serum from patients 1, 2, 3, and 5 during periods with small ovaries and periods with large ovarian cysts (Table 3). In this bioassay, normal pubertal girls had serum FSH bioactivity of 32 ± 4 ng/ml ($n = 6$). Normal adult women in early follicular phase of the menstrual cycle had FSH bioactivity of 202 ± 71 ng/ml ($n = 6$). We have not been able to detect FSH bioactivity in normal prepubertal subjects to date. In contrast, two girls with McCune-Albright syndrome (patients 1 and 5) had detectable FSH bioactivity despite low levels of FSH by radioimmunoassay. Patient 3 had FSH bioactivity in one sample that was at the borderline of detectability. Unilateral ovarian enlargement was evident by ultrasound at the time FSH bioactivity was increased in patients 1, 3, and 5. However, no clearly evident increase in FSH bioactivity was seen in patient 2 at times when large ovarian cysts were present in her ovaries.

DISCUSSION

Girls with McCune-Albright syndrome may have a number of endocrinological abnormalities in addition to sexual precocity. Those include hyperfunction of the thyroid (24), Cushing syndrome (25), and pituitary abnormalities (22). The cause of the endocrine hyperfunction is, as yet, unknown. Hall and Warrick (26) proposed the etiology was related to abnormal or excessive signals from the hypothalamus. However, the majority of patients with McCune-Albright syndrome and sexual precocity have very low levels of gonadotropins (3, 4, 6) and do not exhibit further suppression of gonadotropins nor cessation of menses when treated with long-acting agonists of LHRH. Thus, excess neurotransmitters or excess LHRH secretion cannot account for sexual precocity in most girls with McCune-Albright syndrome, and other etiologies must be investigated.

We have documented that all six girls in our study who had gonadotropin-independent precocious puberty also had enlarged ovaries. Five of the six had intermittent presence of large unilateral ovarian cysts. Ovarian growth with cyst formation appears to be responsible for increases in estradiol production in these girls. Surgical examination has shown that these cysts are generally follicular (2). Despite the formation of such cysts, we and others have not demonstrated elevated progesterone levels in any patients indicating that ovulation has not occurred (2) (Foster CM, unpublished observations).

Ovarian cysts have been seen in association with precocious puberty. Notably, girls with severe hypothyroidism may develop large, multicystic ovaries (21) in response to hormonal overlap from TSH or prolactin (27). TSH shares a common α -subunit with the gonadotropins. However, none of our patients had high TSH levels. Prolactin levels were normal in four of the five girls tested with gonadotropin-independent precocious puberty. This makes an overlap in hormonal sensitivity less likely in these patients.

Circulating factors with gonadotropin bioactivity but little gonadotropin immunoactivity theoretically could stimulate ovarian growth in our patients. Previously we determined LH bioactivity using the rat interstitial cell testosterone production assay in patients 1 and 3 to 6 (3). LH bioactivity was not detectable in any of the girls with low gonadotropin levels. This is in agreement with others who have determined LH bioactivity in girls with McCune-Albright syndrome (6) (Beitins IZ, unpublished observation). In contrast, FSH bioactivity was detected in at least two of our patients. FSH bioactivity did not correlate well with ovarian cyst presence nor with estradiol level. Thus, the significance of this observation is, as yet, unclear.

The fluctuation of ovarian cyst size with consequent fluctuation of plasma estradiol levels appears to be responsible for sexual precocity in girls with McCune-Albright syndrome. Each girl appears to have her own rhythm of cyst production. Cysts have appeared monthly or bimonthly in some of our patients whereas others had cyst formation and exacerbation of symptoms after prolonged quiescent periods of a year or more. We and others have observed that removal of the cyst results in only temporary improvement as remaining ovarian tissue may form cysts (8). Hence surgical intervention is not usually beneficial. Experimental therapy based on interference with estradiol production by use of aromatase inhibitors such as Δ^1 -testolactone has shown more promise (28). The decrease in ovarian cyst size with testolactone therapy suggests that, possibly, there is an imbalance between estrogens and androgens which allows an ovarian cyst to grow. Improved understanding of the cause of ovarian growth in girls with McCune-Albright syndrome will be necessary to improve treatment and outcome.

REFERENCES

1. Albright F, Butler AM, Hampton AC, Smith PH 1937 Syndrome characterized by osteitis fibrosa disseminata, areas of pigmentation and endocrine dys-

- function. *N Engl J Med* 216:727-746
2. Benedict P 1966 Endocrine features in Albright's syndrome. *Metabolism* 11:30-45
 3. Foster CM, Ross JL, Shawker T, Pescovitz OH, Loriaux DL, Cutler GB Jr 1984 Absence of pubertal gonadotropin secretion in girls with McCune-Albright syndrome. *J Clin Endocrinol Metab* 5:1161-1165
 4. Comite F, Shawker TH, Pescovitz OH, Loriaux DL, Cutler GB Jr 1984 Cyclical ovarian function resistant to treatment with an analogue of luteinizing hormone releasing hormone in McCune-Albright syndrome. *N Engl J Med* 311:1032-1036
 5. Foster CM, Comite F, Pescovitz OH, Ross JL, Loriaux DL, Cutler GB Jr 1984 Variable response to a long-acting agonist of luteinizing hormone-releasing hormone in girls with McCune-Albright syndrome. *J Clin Endocrinol Metab* 59:801-805
 6. Wierman ME, Beardsworth DE, Mansfield M, Badger TM, Crawford JD, Criesler JF Jr, Bode HH, Loughlin JS, Kushner DC, Scully RE, Hoffman WH, Crowley WF Jr 1985 Puberty without gonadotropins. A unique mechanism of sexual development. *N Engl J Med* 312:65-72
 7. Pray LG 1951 Sexual Precocity in females report of two cases with arrest of precocity in the McCune-Albright syndrome after removal of a cystic ovary. *Pediatrics* 8:684-692
 8. Case records of the Massachusetts General Hospital (case 4-1975) 1975 *N Engl J Med* 292:199-203
 9. Tanner JM 1978 *Growth at Adolescence*. Blackwell, Oxford
 10. Comite F, Cutler GB Jr, Rivier J, Vale W, Loriaux DL, Crowley WF Jr 1981 Short term treatment of idiopathic precocious puberty with a long-acting analogue of luteinizing hormone releasing hormone. *N Engl J Med* 305:1546-1550
 11. Mansfield MJ, Beardsworth DE, Loughlin JS, Crawford JD, Bode HH, Rivier J, Vale W, Kushner DC, Criegler JF Jr, Crowley WF Jr 1983 Long term treatment of central precocious puberty with a long-acting analogue of luteinizing hormone-releasing hormone Effects on somatic growth. *N Engl J Med* 309:1286-1290
 12. Cargille CM, Rayford PL 1970 Characterization of antiserum for human follicle stimulating hormone radioimmunoassay. *J Lab Clin Med* 75:1030-1040
 13. Odell WD, Ross GT, Rayford PL 1967 Radioimmunoassay for luteinizing hormone in human plasma or serum physiological studies. *J Clin Invest* 46:248-255
 14. Loriaux DL, Ruder HT, Lipsett MB 1971 The measurement of estrone sulfate in plasma. *Steroids* 18:463-472
 15. Aubert AL, Becker RL, Saxina AA, Raiti S 1974 Report of the National Pituitary Agency. Collaborative study of the radioimmunoassay of human prolactin. *J Clin Endocrinol Metab* 38:1115-1120
 16. Ridgeway EC, Weintraub BD, Ceballos JL, Rack MC, Maloof FF 1973 Suppression of pituitary TSH secretion in the patient with hypersecreting thyroid modules. *J Clin Invest* 52:2783
 17. Rich K, Bardin CW, Gunsalus G, Mather J 1983 Age-dependent pattern of androgen binding protein secretion from rat Sertoli cells in primary culture. *Endocrinology* 113:2284-2293
 18. England BG, Niswender GD, Midgley AR 1974 Radioimmunoassay of estradiol 17 beta without Chromatography. *J Clin Endocrinol Metab* 38:42-50
 19. Padmanabhan V, Chappel SC, Beitins IZ 1985 An improved in vitro FSH bioassay suitable for the measurement of hFSH in serum. The Endocrine Society Program and Abstracts of the 67th Annual Meeting; 1136A
 20. Sample W, Dippe BM, Gypes MT 1977 Gray scale ultrasonography of the normal female pelvis. *Radiology* 125:477-483
 21. Lindsay AN, Voorhess ML, MacGillivray MH 1980 Multicystic ovaries detected by sonography in children with hypothyroidism. *Am J Dis Child* 134:589-592
 22. Lightner ES, Penny R, Frasier SD 1975 Growth hormone excess and sexual precocity in polyostotic fibrous dysplasia (McCune-Albright syndrome): evidence for abnormal hypothalamic function. *J Obstet Gynecol* 87:922-927
 23. Carr D, Mathie K, Manners AR, Colman C 1975 *Br J Obstet Gynecol* 86:330-331
 24. Hamilton CR, Maloof F 1973 Unusual types of hyperthyroidism. *Medicine* 52:195-215
 25. Danon M, Robboy S, Kim S, Scully R, Crawford JD 1975 Cushing syndrome, sexual precocity, and polyostotic fibrous dysplasia of bone (Albright syndrome) in infancy. *J Pediatr* 87:917-921
 26. Hall R, Warrick C 1972 Hypersecretion of hypothalamic releasing hormones. A possible explanation of the endocrine manifestations of polyostotic fibrous dysplasia (Albright syndrome). *Lancet* 1:1313-1316
 27. VanWyk JJ, Grumbach MM 1960 Syndrome of precocious menstruation and galactorrhea in juvenile hypothyroidism: an example of hormonal overlap in pituitary feedback. *J Pediatr* 57:416-435
 28. Foster CM, Pescovitz OH, Comite F, Shawker T, Loriaux DL, Cutler GB Jr 1985 Testolactone treatment of precocious puberty due to McCune-Albright syndrome. *Acta Endocrinol* 109:254-257
 29. Shawker TH, Comite F, Rieth KG, Dwyer AJ, Cutler GB Jr, Loriaux DL 1984 Ultrasound evaluation of female isosexual precocious puberty. *J Ultrasound Med* 3:309-316